Quantum phases and phase transitions of interacting bosons in a Kagome lattice

Jinwu Ye
Department of Physics, The Pennsylvania State University, University Park, PA, 16802
(Dated: September 11, 2008)

By using the dual vortex method and analyzing the saddle point structures of the dual gauge fields self-consistently in the insulating side, we study quantum phases, especially supersolid phases and phase transitions in an extended boson Hubbard model at and slightly away from 1/3 (2/3) filling on a kagome lattice. We find phases and phase transitions in Kagome lattice are dramatically different from those in bipartite and triangular lattices in both Ising and Easy-plane limit. Slightly away from $f = 1/3(2/3)$, there is no supersolid in the Ising limit, but there are always supersolids in the Easy-plane limit where we find a stripe solid and a new type of supersolid which has charge density wave-valence bond-superfluid order and also analyze their corresponding stability conditions. Implications on QMC simulations with both nearest neighbor and next nearest neighbor interactions are given.

arXiv:0804.3429v2 [cond-mat.str-el] 11 Sep 2008

1. Introduction. The Extended Boson Hubbard model (EBHM) with various kinds of interactions, at various kinds lattices (bi-partisan or frustrated) at various kinds of filling factors (commensurate or incommensurate) is described by the following Hamiltonian:

$$H = -t\sum_{<ij>}(b_i^\dagger b_j + h.c.) - \mu\sum_i n_i + \frac{U}{2}\sum_i n_i(n_i-1) + V_1\sum_{<ij>} n_i n_j + V_2\sum_{<<ik>>} n_i n_k + \cdots \quad (1)$$

where $n_i = b_i^\dagger b_i$ is the boson density, t is the hopping amplitude, U, V_1, V_2 are onsite, nearest neighbor (nn) and next nearest neighbor (nnn) interactions between the bosons. The $\cdots$ may include further neighbor interactions and possible ring-exchange interactions. A supersolid in Eqn.1 is defined as to have both off-diagonal long range order $< b_i > \neq 0$ and diagonal charge density wave in the boson density n_i. Possible supersolid states in frustrated lattices such as triangular and Kagome lattices were first investigated in [1] by spin wave expansion. A Kagome lattice (Fig.1a) has 3 sublattices a, b and c, it contains both hexagon and triangle, its dual lattice is the dice lattice (Fig.2b). From spin wave expansion, the authors in [1] found that a SS state is more robust in a triangular lattice with only t and V_1 terms in Eqn.1 and a SS state with $\sqrt{3}\times\sqrt{3}$ pattern is stable even at half filling $f = 1/2$. Indeed, this discovery was confirmed by several recent QMC simulations [2]. However, due to too strong quantum fluctuations, the authors in [1] found that the spin wave expansion does not work anymore in Kagome lattice. So a completely different analytical method is needed to study possible supersolid phases in a Kagome lattice.

Recently, the EBHM in square lattice at generic commensurate filling factors $f = p/q$ (p, q are relative prime numbers) were systematically studied in [3] by a dual vortex method (DVM). However, it was explicitly pointed out in [5] that the DVM developed in [3] holds only in the featureless superfluid (SF) and the valence bond solid (VBS) side where the saddle point of the dual gauge field can be taken as uniform, but it fails in the charge density wave (CDW) side where the saddle point of the dual gauge field can *not* be taken as uniform anymore. So the DVM developed in [3] is not self-consistent in the CDW side, the CDW order parameter constructed in [3] to characterize the symmetry breaking patterns in the CDW side are also not self-consistent. So special care is needed to choose a correct saddle point of the dual gauge field in the CDW side, so a different effective action is needed in the CDW side to make the theory self-consistent [5]. In [5], (1) by extending the DVM explicitly to the lattice symmetry breaking side by choosing the corresponding self-consistent saddle points of the dual gauge field (2) by pushing the DVM to slightly away from commensurate filling factors, the author mapped out the global phase diagram of the EBHM on bipartite lattices such as honeycome and square lattice near half filling and investigate superfluid, solid, especially supersolids and quantum phase transitions in the global phase diagram in a *unified* scheme.

In this paper, I will use the DVM developed in [3, 4, 5] to study quantum phases and phase transitions in the kagome lattice at and near filling factor $f = 1/3$ (Fig.2). In a frustrated lattice, the sign of the hopping t in Eqn.1 makes crucial differences. It is also well known that the physics in Kagome lattice is much more complicated than that in a triangular lattice as indicated by the spin wave calculations in [1]. We will also compare our results with some previous Quantum Monte-Carlo (QMC) results in the Kagome lattice [8, 9] and also give important implications to possible future QMC simulations on Eqn.1 with both hard core and soft core, with both the *nn* V_1 and the *nnn* V_2 interactions.

2. The effective action. Because of the P-H symmetry at $U = \infty$, the results are equally valid near 2/3 filling. As shown in [4] at $q = 2$, because the lowest dual vortex band in the dice lattice is completely flat, so the dual vortices are completely localized, so it has to be a super-

fluid, in sharp contrast to $q = 2$ on triangular lattice [6]. For $q = 3$, from Table 1 of [4], we can see there are two minima at $(0, 0)$ and $(-2\pi/3, 2\pi/3)$. Let's label the two eigenmodes at the two minima as $\psi_l, l = 0, 1$. The effective action invariant under all the MSG transformations upto sixth order terms was written down in [8]. Again, inside the superfluid phase, moving *slightly* away from the 1/3 filling $f = 1/3$ corresponds to adding a small *mean* dual magnetic field $\delta f = f - 1/3$ in the action derived in [8]. Upto sixth order, the action is $\mathcal{L}_{SF} = \mathcal{L}_0 + \mathcal{L}_1 + \mathcal{L}_2$:

$$\begin{aligned}
\mathcal{L}_0 &= \sum_l |(\partial_\mu - iA_\mu)\psi_l|^2 + r|\psi_l|^2 \\
&+ \frac{1}{4}(\epsilon_{\mu\nu\lambda}\partial_\nu A_\lambda - 2\pi\delta f \delta_{\mu\tau})^2 \\
\mathcal{L}_1 &= u(|\psi_0|^2 + |\psi_1|^2)^2 - v(|\psi_0|^2 - |\psi_1|^2)^2 \\
\mathcal{L}_2 &= w[(\psi_0^* \psi_1)^3 + h.c.] \qquad (2)
\end{aligned}$$

where A_μ is a non-compact $U(1)$ gauge field.

In fact, the form of this action is *exactly the same* as that in honeycomb lattice at $q = 2$ first derived in [5]. However, the physical meanings of the two eigenmodes $\psi_l, l = 0, 1$ are completely different in the two lattices, this fact leads to completely different physics in the two lattices. In the following, we focus on the insulating side where $r < 0$.

3. Ising limit. If $v > 0$, the system is in the Ising limit, only one of the 2 vortex fields condenses. Let us take $\langle\psi_0\rangle \neq 0, \langle\psi_1\rangle = 0$. It was found that there is no CDW order in [8], but there is a VBS order around either up or down *triangles* (Fig.1). This is just *opposite* to bipartite and triangular lattices discussed in [2, 5, 6, 7] where the Ising limit corresponds to the CDW case. We interpret $\psi_l, l = 0, 1$ as the kinetic energies around the up and down triangles $K_{ijk} = -t(b_i^\dagger b_j + b_j^\dagger b_k + b_k^\dagger b_i + h.c.)$ where i, j, k are the three vertices of a triangle, namely, themselves are valence bond order parameters, while in bipartite and triangular lattices, they are CDW order parameters. This interpretation is indeed consistent with the phases to be identified in the Easy-plane limit in the following. Because the saddle point structure stays the same across the SF to the triangle-VBS (TVB) transition at $f = 1/3$, Eqn.2 still holds in the TVB side, so the transition is a *weak* first order one which breaks both the $U(1)$ and the Z_2 exchange symmetry between ψ_0 and ψ_1[12]. In contrast, in bipartite and triangular lattices, in the Ising limit, different saddle points for the dual gauge fields need to be chosen [5, 6]. These crucial differences are responsible for the absence of supersolid in the Ising limit in a Kagome lattice as shown in the Fig.1. Slightly away from 1/3 filling, there must be a direct frist order transition from the TVB to the superfluid, there is no TVB supersolid intervening between the TVB and the SF in the Fig.1, in sharp contrast to the cases in bipartite and triangular lattices.

4. Easy-plane limit. If $v < 0$, the system is in the easy-plane limit, Eqn.2 is similar to the action in Bilayer quantum Hall systems [11], the two vortex fields have equal magnitude $\psi_l = |\psi|e^{i\theta_l}$ and condense. The relative phases can be determined by the sign of w[5]. In this case, the system is expected to have either CDW or both CDW and VBS around a *hexagon*. As explained in the introduction, if there is any CDW component, Eqn.2 does not work anymore, so can not be used to characterize the symmetry breaking pattern in the insulating side. However, the results in [8] can still be used as a useful guide to see some signatures of the symmetry breaking patterns. In the following, we discuss both cases separately.

(a) Both CDW and VBS order: The analysis in [8] suggests that this is the case when $w > 0$, so $\theta_- = \theta_0 - \theta_1 = (2n+1)\pi/3$ in Eqn.2. Eqn.1 in Kagome lattice with $U = \infty, V_1 > 0$ was studied by recent QMC simula-




FIG. 1: (a) The triangular valence bond (TVB) in a Kagome lattice at $f = 1/3$. The thick triangle means one boson is moving around the up triangle (b) The phase diagram driven by the chemical potential μ. There is a direct first order transition from the TVB to the superfluid. There is no supersolid intervening between the TVB and the SF.




FIG. 2: (a) A Kagome lattice has 3 sublattices labeled as a, b, c. The CDW+VBS in a Kagome lattice at $f = 2/3$ identified in [9]. The dot means one (or $1 - \alpha$) boson occupying the site. The ◯ inside the hexagon means 3 (or $3(1 + \alpha)$ bosons hopping around the hexagon. (b) The saddle point of the dual gauge field in a magnetic field unit cell in the dice lattice.

tions [9, 10]. It was found that the solid state at exactly 1/3 filling has both CDW and VB order. The VB order corresponds to boson hopping around a hexagon (shown in Fig.2a). This is a unique feature of the Kagome lattices. This feature can be intuitively understood without any calculation: with very large V_1 interaction at 1/3 filling, if a boson taking sublattice a, then all its 4 nn which belong to sublattices b and c are excluded, so bosons will go to the 4 nnn which still belong to b and c instead of a (Fig.2a). This is in sharp contrast to triangular lattice at 1/3 filling in [6]. So bosons may move around the hexagon to form some VB order in addition to the CDW order (Fig.2a). Indeed, the up and down triangles in the Fig.2a, each containing one CDW site and one VB, are equivalent. The saddle point for the dual gauge fields is $< \nabla \times \vec{A}_{CDW} >= 1-\alpha, < \nabla \times \vec{A}_{VBS} >= (1+\alpha)/2$ as shown in Fig.2b. The lowest energy dual vortex band in such a saddle point is found to be: $E(k_1,k_2) = -\sqrt{6+2A(k_1,k_2)}$ where $A(k_1,k_2) = \cos k_1 - \cos(k_1-\alpha\pi) + \cos k_2 - \cos(k_2+\alpha\pi) + \cos(k_1-k_2) - \cos(k_1-k_2-2\alpha\pi), -\pi < k_1, k_2 < \pi$. It is easy to see that there is a symmetry $E(k_1,k_2) = E(k_2+\pi(1+\alpha), k_1+\pi(1-\alpha))$. For $0<\alpha<0.404$, there are two minima at $k_1 = \alpha\pi/2 - arcsin(\frac{\sin(\alpha\pi/2)}{2\cos(\alpha\pi)}), k_2 = -k_1$ and $k_1 = \alpha\pi/2 + arcsin(\frac{\sin(\alpha\pi/2)}{2\cos(\alpha\pi)}) + \pi, k_2 = -k_1$. The two minima are related by the symmetry.

When $\alpha = 1/3 < 0.404$, it reduces to $q=3$ case where there are two minima at $(0,0), (-\frac{2\pi}{3}, \frac{2\pi}{3})$ which is the two minima at the superfluid side in Eqn.2. For $0 < \alpha < 1/3^-$, let's still label the two minima by $\psi_l, l = 0, 1$. The $\alpha \to 1/3^-$ case corresponds to approaching to the transition from the CDW+VB to the SF in Fig.3a from the left hand side. Then the MSG inside the SF state $T_1, T_2, R_{\pi/3}, I_1$ is reduced to $T_1, T_2, R_\pi, I_{\vec{a}_1+\vec{a}_2}$ inside the CDW+VBS state in Fig.2b. The effective action slightly away from 1/3 invariant under the MSG transformations inside the CDW+VBS state upto sixth order terms is $\mathcal{L}_{CDW+VBS} = \tilde{\mathcal{L}}_0 + \tilde{\mathcal{L}}_1 + \tilde{\mathcal{L}}_2$:

$$\begin{aligned}
\tilde{\mathcal{L}}_0 &= \sum_l |(\partial_\mu - iA^{VBS}_\mu)\psi_l|^2 + \tilde{r}|\psi_l|^2 \\
&+ \frac{1}{4}(\epsilon_{\mu\nu\lambda}\partial_\nu A^{VBS}_\lambda - 2\pi\delta f \delta_{\mu\tau})^2 \\
\tilde{\mathcal{L}}_1 &= \tilde{u}(|\psi_0|^2 + |\psi_1|^2)^2 - \tilde{v}(|\psi_0|^2 - |\psi_1|^2)^2 \\
\tilde{\mathcal{L}}_2 &= \tilde{w}[(\psi_0^* \psi_1)^2 + h.c.] \qquad (3)
\end{aligned}$$

where A^{VBS}_μ is a non-compact $U(1)$ gauge field. Note that A^{CDW}_μ is always massive, so was already integrated out in Eqn.3. So the density fluctuations on the dot sites in Fig.2a are suppressed. Because the system is already in the CDW+VBS, so we expect $\tilde{r} < 0$ and $\tilde{v} < 0$. Then we can set $\psi_l = |\psi| e^{i\theta_l}$. The relative phases can be determined by the sign of $\tilde{w} > 0$ which has the same sign as $w > 0$. It is easy to observe that due to the same structure of the saddle points on the SF side Eqn.2 and on the CDW-VBS side Eqn.3, the SF to the CDW+VB transition driven by r along the horizontal axis in Fig.3a is very weakly first order. This is in sharp contrast to the strong 1st order SF to CDW transition in a triangular lattice at $f = 1/3$ studied in [6]. It is also different from the SF to the VBS transition at $f = 1/2$ in honeycomb and square lattices where the same action can be used on both side. Although the QMC in Ref.[10] found the CDW-VB solid to the SF at $f = 1/3$ is a 2nd order transition through a possible deconfined quantum critical point, the QMC in Ref.[9] found a weakly 1st order transition from double-peaked histogram of the boson kinetic energy. Our results from the self-consistent DVM explains the underlying mechanism of the weak first order transition.

When moving from 1/3 to 1/2 where the boson has to be a SF, Eqn.3 can be rewritten as:

$$\begin{aligned}
\tilde{\mathcal{L}} &= (\frac{1}{2}\partial_\mu \theta_+ - A^{VBS}_\mu)^2 + \frac{1}{4e^2}(\epsilon_{\mu\nu\lambda}\partial_\nu A^{VBS}_\lambda - 2\pi\delta f\delta_{\mu\tau})^2 \\
&+ (\frac{1}{2}\partial_\mu\theta_-)^2 + 2\tilde{w}\cos 2\theta_- \qquad (4)
\end{aligned}$$

where $\theta_\pm = \theta_0 \pm \theta_1$.

The SS slightly away from 1/3 filling has the same CDW and VB order as the solid at 1/3, so let's call this kind of novel SS as CDW-VB-SS. Then the transition from the CDW+VB to the CDW-VB-SS transition driven by the chemical potential in Eqn.4 can be similarly discussed as the VBS to the VB-SS in bipartite lattices in [5, 6, 7], so it is is the same universality class as that from the Mott insulator to the SF. The CDW-VB-SS to the SF transition is 1st order. Although the CDW-VB-SS may not be stable in the hard core case, it should be stable in the soft core case. Of course, there is no P-H symmetry anymore in the soft core case. This study may inspire more accurate QMC to search for this novel CDV-VB-SS state, especially in the soft-core case.

(b) Only CDW order: We expect that this is the case when $w < 0$, so $\theta_- = \theta_0 - \theta_1 = 2n\pi/3$ in Eqn.2. If we add very strong *nnn* interaction V_2, the bosons will go to the *nnnn* which belong to a again (Fig.2a), then the bosons will simply take one of the 3 sublattices such as a to form a CDW alone (Fig.3b). Indeed, the up and down triangles in the Fig.2a, each containing one occupied site and two empty sites, are equivalent. In the stripe CDW state in Fig.3b, a different saddle point where $< \nabla \times \vec{A}^a >= 1-2\alpha$ for the sublattice a and $< \nabla \times \vec{A}^{bc} >= \alpha$ for the two sublattices b and c should be used. The $\alpha \to \alpha_c < 1/3$ limit corresponds to approaching the stripe CDW to the SF transition from the stripe CDW side in Fig.3b, while the $\alpha \to 0^+$ limit corresponds to $t/V_1 \to 0$ limit in Fig.3b. It is easy to see that there is only one vortex minimum ψ_{bc} in such a staggered dual magnetic field with $\alpha < \alpha_c < 1/3$, so the effective action




FIG. 3: (a) Zero temperature phase diagram in Kagome lattice $U = \infty, V_1 > 0$. In the soft core case $U < \infty$, there maybe a narrow window of CDW-VB-SS sandwiched between the CDW-VB solid and the SF. In the hard core case studied in [9, 10], the CDW-VB-SS is unstable, there is a direct weakly first order transition from the CDW-VB solid to the SF. (b) Very similar phase diagram also works for $U = \infty, V_1 > 0, V_2 > 0$. In this case, V_1 in (a) need to be replaced by $V_1 + V_2$, the solid at 1/3(2/3) is simply a CDW where the bosons occupying (un-occupying) one of the 3 sublattices a, b, c in Fig.2a. The SS is simply a stripe SS where the superfluid stiffness $\rho_{bc} \gg \rho_{ba}$. The Stripe-SS will surely be stable even in the hard core case. As shown in the text, the first order transition in (a) is a weakly first order one, but strongly first order one in (b). Thin (thick) line is 2nd (first) order transition.

inside the stripe CDW state is:

$$\mathcal{L}_{stripe} = |(\partial_\mu - iA^{bc}_\mu)\psi_{bc}|^2 + r_s|\psi_{bc}|^2 + u_s|\psi_{bc}|^4 + \cdots + \frac{1}{4}(\epsilon_{\mu\nu\lambda}\partial_\nu A^{bc}_\lambda - 2\pi\delta f \delta_{\mu\tau})^2 \quad (5)$$

where the vortices in the phase winding of ψ_{bc} should be interpreted as the the boson number. Because of sharp change of the saddle point from Eqn.2 in the SF side to Eqn.5 in the stripe CDW side, the transition from the SF to the stripe CDW along the horizontal axis in Fig.3b is likely to be a strong first order.

Slightly away from 1/3, there is a stripe SS as shown in Fig.3b. In the direct lattice Fig.2a, even if taking away the sublattice a, the bosons can still move easily along the chain *bcbc...*, so they need only overcome the barrier along the chain *baba...* to achieve an effective hopping along *baba....* So the superfluid stiffness along the *bc* chain is larger than that along the *ba* chain. Because the *nn* and the *nnn* neighbors are similar as can be seen from Fig.2a, the vertical axis in Fig.3b need to be replaced by $\mu/(V_1 + V_2)$, the horizontal axis stay as t/V_1 at fixed large V_2 (Fig.3b). While the conclusion achieved in [4] that the system has to be a superfluid state at 1/2 remains robust. So we conclude that when moving from 1/3 to 1/2 driven by the chemical potential along the vertical axis in Fig.3b, the system goes CDW to stripe-SS to the SF transition. The first is the second order, the second is a first order transition (Fig.3b). Again, longer-range interactions favor the description in terms of the dual vortices and the stability of SS, so the Stripe-SS will surely be stable even in the hard core case (Fig.3b). We suggest that it is important to do QMC with $U = \infty, V_1 > 0, V_2 > 0$ to test Fig.3b.

5. Conclusions. By using the DVM developed in [3, 5, 6], we studied quantum phases and phase transitions in an extended boson Hubbard model at and slightly away from 1/3 (2/3) filling on a kagome lattice. We find that in the insulating side, at the commensurate fillings such as 1/3, one has to choose the corresponding saddle point structure of the dual gauge fields self-consistently to construct effective actions on this side. We find quantum phases and phase transitions in the Kagome lattice in both Ising and easy plane limit are dramatically different from those in bipartite and triangular lattice studied previously [2, 3, 5, 6, 7]. We interpret the two vortex order parameters $\psi_l, l = 0, 1$ as the kinetic energies around the up and down triangles in a Kagome lattice. In the Ising limit, the SF to the triangle valence bond order (TVB) is a weak first order one. For the first time, the Ising limit of the same action can be used to describe a quantum phase transition in a concrete model. In the easy-plane limit, we find that the CDW+VB to the SF transition at commensurate filling $f = 1/3$ in Kagome lattice is a weakly first order one. This finding resolves some outstanding puzzles observed in the QMC in a kagome lattice [9, 10]. When pushing the effective action slightly away from the commensurate fillings, we find that in the Ising limit, there can only be a direct first-order transition from the triangular valence bond (TVB) to the SF, so there is no immediate TVB supersolid intervening between the TVB and the SF. However, in the Easy-plane limit, different kinds of supersolids are generic stable states slightly away from 1/3 (2/3) filling. In addition to a stripe supersolid (stripe-SS), we also find a new kind of supersolid: CDW-VB supersolid. The stripe-SS should be stable even in the hard core limit, while the CDW-VB-SS maybe stable only in the soft core limit. We also made implications on the available and future QMC simulations on EBHM with both *nn* and *nnn* interactions in a kagome lattices.

I am indebted to Longhua Jiang for technical assistance in finding the saddle point structures in the Kagome lattice.

[1] Ganpathy Murthy, Daniel Arovas, Assa Auerbach, Phys. Rev. B 55, 3104 (1997).
[2] S. Wessel, M. Troyer, Phys. Rev. Lett. 95, 127205 (2005); M. Boninsegni, N. Prokof'ev, *ibid* 95, 237204 (2005).
[3] Balents L, Bartosch L, Burkov A, Sachdev S, Sengupta K, Phy. Rev. B 71, 144508 (2005).
[4] Longhua Jiang and Jinwu Ye, J. Phys. Condens. Matter 18, 6907-6922 (2006).

[5] Jinwu Ye, cond-mat/0503113
[6] Jinwu Ye, arXiv:0712.0251, to be published in Nucl.Phys. B.
[7] Jing Yu Gan, Yu Chuan Wen, Jinwu Ye, Tao Li, Shi-Jie Yang, Yue Yu, Phys. Rev. B 75, 214509 (2007).
[8] K. Sengupta, S. V. Isakov, Yong Baek Kim, Phys. Rev. B 73, 245103 (2006).
[9] S. V. Isakov, S. Wessel, R. G. Melko, K. Sengupta, Yong Baek Kim, Phys. Rev. Lett. 97, 147202 (2006).
[10] K. Damle, T. Senthil, Phys. Rev. Lett. 97, 067202 (2006).
[11] Jinwu Ye and Longhua Jiang, Phys. Rev. Lett. 98, 236802 (2007); Jinwu Ye, Phys. Rev. Lett. 97, 236803 (2006)
[12] For quantum phase transitions in Ising limit in the contect of bilayer quantum Hall systems in a periodic potential, see appendix C in Jinwu Ye, Annals of Physics, 323 (2008), 580-630.